

Dear Dr. Farke,

Please find attached the revised version of our manuscript entitled “The impact of allometry on vomer shape and its implications for the taxonomy and cranial kinesis of crown-group birds”.

Both reviewers provided useful comments, and we are happy to acknowledge that we included most of their suggestions to the new version. In the few cases where we did not agree with the reviewer, we provided a detailed response. All responses to the reviewers are highlighted in bold.

We hope that you are satisfied with the revised version and looking forward to hear from you.

With kind regards,

Christian Foth (on behalf of all authors)

Reviewer#1

Present study lead by Plateau & Foth and titled ‘*The impact of allometry on vomer shape and its implications for the taxonomy and cranial kinesis of crown-group birds*’ deeply addresses a relevant aspect that strongly affect the morphological evolution of several anatomical structures: size. The approach you propose is essential to correctly understand how the influence of this central structure is in craniofacial and PPC configuration, and also in kinesis performance among other cranial traits. This is important specially when such structures are suggested as taxonomic distinguishing. Thus, it is important that you emphasize that, although there are important morphological differences between the main groups of modern birds, these are mostly due to (or related to) the evolution of body size, a fundamental biological trait, and not so much as shape changes themselves (i.e. vomer morphology is not independent, in this case from its size, and if you don’t take raw size differences as a taxonomic trait, you shouldn’t take as such a morphological trait associated with it).

The authors address this important question with a wide and dense combination of methods which look for the pure morphological differences that are not under the widespread influence of size, but I suggest below some commonplace aspects that I think they could contribute to the robustness of the study and make it more complete.

- **We are thankful for the overall positive evaluation of our study.**

Methods

I would suggest to estimate the morphological disparity from the Procrustes shape variables for each group and compare it between groups with pairwise comparisons. Thus, you can compare statistically the morphological variation for each group and their distinct distribution across the morphospace, especially when you extract size and groups collapse, obtaining different significant results in size-affected shapes.

- **We included a disparity analysis showing the morphospace size of different birds groups and how it changes when shape is corrected for allometry. Although disparity analyses allow to test for size, density and position of the morphospace, this method does not allow evaluating the spatial overlap of single morphospaces.**

Also, another option to provide robustness to the hypothesis that vomer's morphology is not a good taxonomic character is estimating the phylogenetic signal both in shape variables, centroid size and centroid size proportional to skull size. You should obtain empirical K values lower to the predicted value of $K = 1$ for a Brownian Motion model of evolution indicating weaker phylogenetic signal and a higher degree of convergence. Additionally, it should be observed a similar distribution across the phylogeny between shape variables and size variables plotting them as continuous variables.

- **We added a short section to the results, showing that shape is affected by phylogeny, while size does not. The method is explained in the method section.**

I would suggest to carry out the allometric regression with Phylogenetic Generalized Least Squares and group by group, running pairwise comparisons between their slopes in order to know how is the allometric influence. Allometry seems to affect specially in Paleognathae considering that when you extract size the difficulty to distinguish these birds from the rest increases. This aspect could be due to ratites reaches bigger sizes.

- **We agree with the reviewer that data should be corrected for phylogeny if possible. Therefore, we applied a phylogenetic flexible discriminate analyses (see line 150ff) and compared it later with the results PERMANOVA and DA results. As the outcome is more or less the same for all types of analyses, the impact of phylogeny is negligible with respect to allometric shape variation.**

Finally, you discuss about kinesis and why is not possible that its different types are related to vomer shape. In order to know this lack of correlation I would recommend grouping the sample by kinesis type, estimating morphological disparities for each group and comparing statistically their distribution across the morphospace. It should collapse widely. Also, it would be interesting to explore by PGLS how

different types of kinesis are related to different allometric scaling, because it could be that the shape-size relationships from distinct kinesis types are similar and not significantly different. Furthermore, to do that I would take into account as an important aspect for kinesis performance the vomer size relative to skull size for a better addressing of their significance on such trait.

- **Apart from Zusi 1984, there is no interspecific study sampling discriminating different types of kinesis in birds with a broad taxon. Unfortunately, Zusi 1984 presented the different types on the family but not species level. Therefore, we cannot apply Zusi's scheme on the sampling of Hu et al. 2019.**

Figures

For the figure 1 I suggest to add a close up of the region that anatomically distinguishes paleognathans from neognathans and a better schedule that could show the stiffness or mobility of the PPC (for example, marking or highlighting the structures that characterize each type of PPC).

- **To highlight the morphological differences between a neognath and palaeognath palate, we highlighted the palatal bones in different colours and indicated the articulation in neognath birds that allow for the kinetic movement of the skull and the connection between braincase and palate in palaeognaths that prevent this kinesis.**

For the PCAs (figure 2) I would suggest trying to plot vomer size and vomer size overall skull size as dot size. Thus, it could be visualized the polarization of size on the morphospace and the wide divergence between paleognathans and neognathans. Finally, the interpretation of the graphics would improve if you add silhouettes representing different groups and if you mark above from which analysis they come or if they are taking size into account or not.

- **We follow the suggestion of the reviewer, adding silhouettes representing the different bird clades and adjusted the dots according log-centroid size of the vomer. We further indicated if the plots sourced on allometric or non-allometric data.**
- **As a note for the editor: We also changed the colours of plots to improve visibility for colour-blind people.**

Reviewer#2

This is an interesting piece of work, and takes further the previous work of Hu et al. (2019), of which a reanalysis of whose data it basically consists. It is a necessary further analysis of the data, and it appears to be sound methodologically and in its basic conclusions. I am not a particular expert in the groups at hand however, and thus my conclusions regarding its soundness are not based on specific knowledge of the taxa.

One major query I have, given the very specific nature of the work, and the fact that it basically consists of a reanalysis of a recent dataset, is perhaps it would be better published in PNAS as a reply to the original paper of Hu et al.? This would give the authors a chance to respond to the comments, and mean it was read by the scientific public as a direct continuation of the original work. ...So yes, perhaps this could be suggested to the authors, if they have not already considered it?

- **Of course, we thought about submitting a response to PNAS. However, these comments are usually limited to not more than 500 words and do not allow the presentation of a sophisticated re-analysis of the data. As we applied several different methods to test the relationship between shape and taxonomy, we decided that it would be better to publish this work separately.**

My only other major comment really, given that the “meat” of the work is pretty sound, is that the discussion ends rather abruptly, and doesn’t expand on the wider implications of the reanalysis of the data of Hu and colleagues. It would be nice to have a concluding paragraph or two at the end of the discussion, with a bit of contextualisation of the study’s findings in terms of their broader implications for avian evolution and evolutionary processes and analysis more broadly. Any main points here could also be added as a sentence of the abstract. ...Specifically, the wider implication discussed by Hu et al. for the radiation of the avian crown group should be addressed, and what differences the reanalysis makes for these conclusions outlined.

- **The reviewer is correct that the discussion of the previous version of the manuscript ended rather abruptly. Therefore, we followed the suggestion of the reviewer and added a conclusion at the end of the discussion. However, we would like to point out that the previous version of the manuscript already discussed implications of reanalysing Hu’s data.**

Some other ideas of wider implications: isolated elements may not reliably group taxa, and can be more strongly subject to convergence; the avian skull has seen a labile pattern of evolution, with repeated development of similar morphology – perhaps this is broadly the case in many groups?; it should be checked that allometry is not the driving cause behind seemingly “primitive” shapes and characteristics etc. etc. Also, can you place the vomer development in birds somehow in the wider context of archosaur evolution, for example of the feeding and sensory systems? There are a few minor linguistic points (mostly just formal v. informal issues, or clarity), which I annotated in the PDF.

- **We already pointed out that the vomer is not a reliable element for taxonomic identifications. However, we have slightly extended this argument in the revised version.**
- **Unfortunately, the vomer is an inconspicuous bone in dinosaurs that has not get much attention so far, including morphology, ecology and evolution. Therefore,**

it is hard to make any meaningful statements for its role in feeding. According to Bertmar (1981, Evolution, 35:359-366), the vomer-associated Jacobson's organ is lost in birds and crocodylians. The situation in non-avian dinosaurs is unknown. So nothing can be said about its sensory role of the vomer.

- **Unfortunately, we did not really understand the reviewer's statement on the role of allometry in terms of primitive shapes. Is he referring to the body size evolution of birds? This statement need to be more specified, before we can address the reviewer's comment.**
- **All other changes suggested in the attached PDF have been implemented.**

In the reference list, the book references seem a bit lacking in details (e.g. number of pages, editors) – if this is acceptable for PCI Palaeo that's OK (ask the editor!), but it is also not consistent throughout currently.

- **For the references, we used the citation style of PeerJ, which does not provide any page numbers for books, only for book chapters. The reviewer might confused the citation of books and book chapters. In the later cases, we provided the authors, chapter title, editors, book title, pages, publisher and location of the publisher. However, we are happy to include also the page numbers for the book references, too.**

In figure captions 2, several abbreviations (e.g. PCA) are used without being defined. I think they should be defined the first time used in each figure caption so that they can be read without referring to the text.

- **This was changed.**

I would recommend adding silhouettes of either whole animals or at least skulls on figures 2 and 3, to make it clearer what is what. The colours are great, but silhouettes would help a lot.

- **As recommended by the reviewer, we included silhouettes to figure 2 and the supplementary figure. We agree that the silhouettes ease the understanding of the taxa-specific morphospaces which are shown as convex hulls. However, we did not do it for figure 3 because we did not use convex hulls and for this kind of plot silhouettes are rather confusing.**

There are also some issues recurring throughout the manuscript with taxon names. These have been annotated the first few times they occur in the PDF, but should be checked throughout:

-The abbreviation of genus names is not consistently done (sometimes not abbreviated) and is confusing. This also goes for figure 1. "S." is used in rapid succession for three different genera! I would recommend you state the full genus name in all cases, and if you want to abbreviate then do so by removing the species name in monospecific taxa. ...If you *really* want to abbreviate genera, can you reduce them by only enough letters so no abbreviations overlap, for example "St." for "Struthio", "Si" for "Sinovenator" etc.?

- **This was changed. All genera names are now written out.**

-The capitalization of Latin versus anglicized clade names does not seem to be always correct. Original Latin (e.g. “Inopinaves”) should be capitalized, and anglicized versions (I presume in this case it would be “inopinavians”) should not. If in doubt, I’d say stick to the Latin and capitalize.

-I guess “aequorlitorithines” is maybe the correct anglicization of “Aequorlitorithes”, but the only place I found it used so far was this preprint! (I guess it’s only been in existence since 2015, so that’s why) Maybe stick to the capitalized Latin? I am not an expert in avian taxonomy though, so your call.

-Sometimes “Galloanseres” or “galloanserinae” have been used

- **All recommended bird groups have been changed to the Latin form.**

-“jackknife” isn’t hyphenated!

- **This has been corrected**