



RECOMMENDATION

Cite as: Jarochovska E (2022). Insights into mechanisms of coccolithophore speciation: How useful is cell size in delineating species? *Peer Community in Paleontology*, 100011. DOI: 10.24072/pci.paleo.100011

Published: 24th October 2022

Based on reviews by:
Andrej Spiridonov and one anonymous reviewer

Correspondence:
e.b.jarochovska@uu.nl

© 2022 Author(s)



CC-BY-ND 4.0

This work is licensed under the Creative Commons Attribution-NoDerivatives 4.0 International License.

Insights into mechanisms of coccolithophore speciation: How useful is cell size in delineating species?

Emilia Jarochovska¹ 

¹ Department of Earth Sciences, Utrecht University – Utrecht, Netherlands

A recommendation of

Razmjooei MJ and Thibault N (2022). Morphometric changes in two Late Cretaceous calcareous nanofossil lineages support diversification fueled by long-term climate change. *PaleorXiv* nfy9, ver. 4, peer-reviewed by PCI Paleo. DOI: 10.31233/osf.io/nfy9

Calcareous plankton gives us perhaps the most complete record of microevolutionary changes in the fossil record (e.g. [Tong et al., 2018](#); [Weinkauff et al., 2019](#)), but this opportunity is not exploited enough, as it requires meticulous work in documenting assemblage-level variation through time. Especially in organisms such as coccolithophores, understanding the meaning of secular trends in morphology warrants an understanding of the functional biology and ecology of these organisms. [Razmjooei and Thibault \(2022\)](#) achieve this in their painstaking analysis of two coccolithophore lineages, *Cribrosphaerella ehrenbergii* and *Microrhabdulus*, in the Late Cretaceous of Iran. They propose two episodes of morphological change. The first one, starting around 76 Ma in the late Campanian, is marked by a sudden shift towards larger sizes of *C. ehrenbergii* and the appearance of a new species *M. zagrosensis* from *M. undulatus*. The second episode around 69 Ma (Maastrichtian) is inferred from a gradual size increase and morphological changes leading to probably anagenetic speciation of *M. sinuosus* n.sp.

The study remarkably analyzed the entire distributions of coccolith length and rod width, rather than focusing on summary statistics ([De Baets et al., In Press](#)). This is important, because the range of variation determines the taxon's evolvability with respect to the considered trait ([Love et al., 2022](#)). As the authors discuss, cell size in photosymbiotic unicellular organisms is not subject to the same constraints that will be familiar to researchers working e.g. on mammals ([Niklas, 1994](#); [Payne et al., 2009](#); [Smith et al., 2016](#)). Furthermore, temporal changes in size alone cannot be interpreted as evolutionary without knowledge of phenotypic plasticity and environmental clines present in the basin ([Aloisi, 2015](#)). The more important is that this study cross-tests size changes with other morphological parameters to examine whether their covariation supports inferred speciation events. The article addresses as well the effects of varying sedimentation rates ([Hohmann, 2021](#)) by, somewhat implicitly, correcting for the stratigraphic trend using an age-depth model and accounting for a hiatus. Such multifaceted approach as applied in this work is fundamental to unlock the dynamics of speciation offered by the microfossil record.

